

Catalog Number: 21188-1, 21188-2

Amount: 50µg/50µl, 100µg/100µl

Swiss-Prot No. : P63104

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl,0.02% sodium azide and 50% glycerol.

Storage/Stability: Store at -20°C/1 year

Immunogen: The antiserum was produced against synthesized non-phosphopeptide derived from

Human 14-3-3 ζ around the phosphorylation site of serine 58 (R-S-S^P-W-R).


Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Specificity/Sensitivity: 14-3-3 ζ (Ab-58) antibody detects endogenous levels of total 14-3-3 ζ protein Reactivity: Human, Mouse, Rat

Applications:

Predicted MW: 28 kd


WB:1:500~1:1000 IHC:1:50~1:100



TNF-α - - +

Peptide - + - -

Western blot analysis of extract from NIH/3T3 cells, untreated or treated with TNF- α (20ng/ml, 5min), using 14-3-3 ζ (Ab-58) antibody (#21188, lane 1 and 2) and 14-3-3 ζ (Phospho-Ser58) antibody (#11181, lane 3 and 4).



Peptide - + Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue, using 14-3-3 ζ (Ab-58) antibody (#21188).

Background :

This gene product belongs to the 14-3-3 family of proteins which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals, and this protein is 99% identical to the mouse, rat and sheep orthologs. The encoded protein interacts with IRS1 protein, suggesting a role in regulating insulin sensitivity. Several transcript variants that differ in the 5' UTR but that encode the same protein have been identified for this gene.

References:

Gu YM, et al. (2006) FEBS Lett ; 580(1): 305-310 Powell DW, et al. (2003) Mol Cell Biol; 23(15): 5376-5387 Mackintosh *C*. (2004) Biochem J; 381(Pt 2): 329-342.